

Clinical decision making: Evolving from the hypothetico-deductive model to knowledge-enhanced machine learning

Han Yuan 

Duke-NUS Medical School, National University of Singapore, Singapore

Correspondence

Han Yuan, Duke-NUS Medical School, National University of Singapore, 8 College Road, Singapore 169857, Singapore.

Email: yuan.han@u.duke.nus.edu

Funding information

None

KEYWORDS

clinical decision making, doctor-in-the-loop, hypothetico-deductive model, machine learning, pattern recognition model

Clinical decision making (CDM) is a process that healthcare professionals undertake when making assessments about patients' conditions and decisions about the care to provide [1, 2]. Traditional CDM is founded on either unconscious intuition or conscious inference frameworks with well-defined logic [3]. Intuition, defined as understanding without a rationale, integrates tacit knowledge and pertinent experience developed over years of practice to automate cognitive processing devoid of formalized rules [4, 5]. However, its unconscious nature obscures the precise identification of initiating cues and inference logic, limiting its application in CDM [6]. Distinct from intuition, conscious inference frameworks possess well-defined logic and execution steps, predominantly encompassing the hypothetico-deductive model (HDM) [7] and pattern recognition model (PRM) [8].

1 | CLASSIC CDM METHODS

The HDM involves four indispensable steps: cue acquisition, hypothesis generation, cue interpretation, and hypothesis evaluation [7]. Initially, cue acquisition

systematically collects patient medical information per requirement by clinicians. Subsequently, multiple preliminary hypotheses are derived from the retrieved information, enabling clinicians to either use an established theory based on gathered information or propose hypotheses according to their clinical knowledge and domain expertise [9]. This is followed by cue interpretation, in which clinicians discern cues pertinent to their initial hypotheses and accordingly refine these hypotheses [10]. The HDM culminates in hypothesis evaluation, whereby the hypotheses are either corroborated or refuted based on the amassed evidence. Should all hypotheses be rejected, another round of the HDM will commence. Ng et al. present an application of the HDM in a real-world CDM task [11]. When diagnosing patients with acute chest pain, clinicians gather cues related to cardiovascular risk factors, smoking history, recent viral infections, and other relevant information. Based on this information, they generate diagnostic hypotheses, such as acute coronary syndrome, myocarditis, pericarditis, or pneumonia. They interpret these cues and evaluate the hypotheses, refining and ruling in or out possibilities through further investigations, such as electrocardiograms, complete blood

Abbreviations: CDM, clinical decision making; HDM, hypothetico-deductive model; KEML, knowledge-enhanced machine learning; ML, machine learning; PRM, pattern recognition model.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2024 The Author(s). *Medicine Advances* published by John Wiley & Sons Ltd on behalf of Tsinghua University Press.

counts, and chest radiographs. In contrast to the analytical HDM, PRM employs nonanalytical matching of new clinical cases with similar patterns stored in memory, specifically for patients encountered previously or documented within established guidelines [8, 10]. In routine clinical encounters, PRM outpaces CDM at an exceptional rate. Notably, in instances of ambiguity, the analytical HDM approach retains its superiority as a more effective solution [8, 10].

2 | CONTEMPORARY CDM METHODS

Although these traditional methods are widely implemented in real-world practice and some of them, including the HDM, have been recognized as gold standards [12], machine learning (ML), as demonstrated in Figure 1, is increasingly adopted to handle the unprecedented volume of information generated by advanced medical instruments and electronically recorded by medical systems [13]. This abundance of data poses challenges for traditional CDM methods relying on manual effort, but ML techniques hold promise because they enable computers to

automatically learn projection functions between raw data and targets of interest without explicit instructions from human experts [14]. For example, support vector machine, random forest, and k-nearest neighbor have been used to diagnose Alzheimer's disease [15], breast cancer [16], and Parkinson's disease [17], respectively. In addition to these conventional techniques, deep learning, a specialized ML subset focusing on design and training strategies of artificial neural networks, has emerged as the state-of-the-art approach for various CDM tasks owing to its extensive parameterization and intricate pattern recognition capability [13].

3 | FUTURE CDM METHODS

Though purely data-driven ML has demonstrated superior accuracy over traditional CDM methods [18], ML exhibits drawbacks in interpretability because of its complex architectures for information processing [19]. Interpretability stands as a pivotal characteristic within CDM to rectify potential erroneous decisions that endanger patient lives. To address this challenge and




FIGURE 1 Comparison of a traditional method (hypothetico-deductive model) versus a contemporary machine learning approach (random forest) in scholarly publications over the last 25 years. Specific numbers were retrieved through a systematic inquiry on Google Scholar employing the search terms “clinical decision making” in conjunction with either “hypothetico-deductive model” or “random forest” on August 21, 2024.

further augment other capabilities of ML, including its exceptional accuracy, researchers collaborate with clinicians to incorporate clinical knowledge and domain expertise into ML methodologies [20]. As depicted in Figure 2, this integration, termed knowledge-enhanced ML (KEML) [21], can be conceptualized as a fusion of clinical knowledge and domain expertise extracted from traditional CDM alongside powerful ML architectures, encompassing both conventional models and deep learning approaches [22].

The foremost advantage of KEML over contemporary ML is its ability to significantly improve interpretability [23]. Although various explainable artificial intelligence techniques have been proposed to supplement ML with interpretability, these data-driven explanations frequently suffer from logical inconsistencies stemming from noise within datasets or limited applicability to particular cohorts [24]. For instance, a previous study showed that data-

driven ML classifiers for pneumothorax often rely on irrelevant regions beyond the lesion area for diagnosis, resulting in overfitting to specific data sources [23]. Integrating clinical knowledge of disease occurrence can enhance both the generalization and interpretability of these classifiers. Another illustrative example is the knowledge-guided interpretable disease prediction method, which showcases the application of medical knowledge graphs to modeling personalized patient data and improving CDM interpretability by extracting crucial graph paths as prompts for ChatGPT to generate clinician-comprehensible natural language explanations [24]. In addition, KEML leverages external clinical knowledge to improve CDM accuracy further. Dynamic gated recurrent neural network exemplifies the enrichment of the representation of a specific medical event with additional information about its adjacent events [25]. KEML also alleviates the biases of data-driven ML through the




FIGURE 2 Schematic plot depicting the fusion of classic and machine learning methods leading toward future knowledge-enhanced machine learning for clinical decision making.

involvement of healthcare professionals and their medical expertise. Chen et al. systematically summarized the potential of KEML to mitigate disparity and inequity in each stage of the ML life cycle [26]. Hence, the integration of clinical knowledge into ML not only amplifies interpretability and accuracy but also mitigates biases, ultimately advancing the real-world deployment of ML-driven healthcare solutions [27].

In this commentary, we summarized the evolution of CDM from traditional to contemporary ML techniques and elucidated the underlying rationale driving this paradigm shift, emphasizing the imperative of adapting to the era of big data. While embracing ML represents a crucial advancement in modeling clinical information, knowledge and expertise embodied in traditional methods should not be disregarded. Therefore, we advocate for KEML, a novel paradigm capitalizing on the strengths of both methodologies, to propel CDM to an exceptional level of interpretability, accuracy, and fairness [28].

AUTHOR CONTRIBUTIONS

Han Yuan: Conceptualization; data curation; formal analysis; investigation; methodology; visualization; writing—original draft; writing—review & editing.

ACKNOWLEDGMENTS

None.

CONFLICT OF INTEREST STATEMENT

The author declares no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no data was generated or analyzed during the current study.

ETHICS STATEMENT

Not applicable.

INFORMED CONSENT

Not applicable.

ORCID

Han Yuan  <https://orcid.org/0000-0002-2674-6068>

REFERENCES

- [1] Banning M. A review of clinical decision making: models and current research. *J Clin Nurs*. 2008;17(2):187–95. <https://doi.org/10.1111/j.1365-2702.2006.01791.x>
- [2] Bright TJ, Wong A, Dhurjati R, Bristow E, Bastian L, Coeytaux RR, et al. Effect of clinical decision-support systems: a systematic review. *Ann Intern Med*. 2012;157(1):29–43. <https://doi.org/10.7326/0003-4819-157-1-201207030-00450>
- [3] Manias E, Aitken R, Dunning T. Decision-making models used by 'graduate nurses' managing patients' medications. *J Adv Nurs*. 2004;47(3):270–8. <https://doi.org/10.1111/j.1365-2648.2004.03091.x>
- [4] Buckingham CD, Adams A. Classifying clinical decision making: a unifying approach. *J Adv Nurs*. 2000;32(4):981–9. <https://doi.org/10.1046/j.1365-2648.2000.t01-1-01565.x>
- [5] Shaban RZ. Theories of clinical judgment and decision-making: a review of the theoretical literature. *Australas J Paramed*. 2005;3:1–13. <https://doi.org/10.33151/ajp.3.1.308>
- [6] Buckingham CD, Adams A. Classifying clinical decision making: interpreting nursing intuition, heuristics and medical diagnosis. *J Adv Nurs*. 2000;32(4):990–8. <https://doi.org/10.1046/j.1365-2648.2000.t01-1-01603.x>
- [7] Elstein AS, Shulman LS, Sprafka SA. Medical problem solving: an analysis of clinical reasoning. Cambridge: Harvard University Press; 1978. <https://doi.org/10.4159/harvard.9780674189089>
- [8] Yazdani S, Hosseinzadeh M, Hosseini F. Models of clinical reasoning with a focus on general practice: a critical review. *J Adv Med Educ Prof*. 2017;5(4):177–84.
- [9] Pearson H. Science and intuition: do both have a place in clinical decision making? *Br J Nurs*. 2013;22(4):212–5. <https://doi.org/10.12968/bjon.2013.22.4.212>
- [10] Woodford P. The decision is yours. *J Paramedic Pract*. 2015;7(2):90–4. <https://doi.org/10.12968/jpar.2015.7.2.90>
- [11] Ng IKS, Goh GW, Teo DB, Chong KM, Tan LF, Teoh CM. Clinical reasoning in real-world practice: a primer for medical trainees and practitioners. *Postgrad Med*. 2024;qgae079. <https://doi.org/10.1093/postmj/qgae079>
- [12] Duong QH, Pham TN, Reynolds L, Yeap Y, Walker S, Lyons K. A scoping review of therapeutic reasoning process research. *Adv Health Sci Educ Theory Pract*. 2023;28(4):1289–310. <https://doi.org/10.1007/s10459-022-10187-7>
- [13] Yuan H, Yu K, Xie F, Liu M, Sun S. Automated machine learning with interpretation: a systematic review of methodologies and applications in healthcare. *Med Adv*. 2024;2(3):205–37. <https://doi.org/10.1002/med4.75>
- [14] Adlung L, Cohen Y, Mor U, Elinav E. Machine learning in clinical decision making. *Med*. 2021;2(6):642–65. <https://doi.org/10.1016/j.medj.2021.04.006>
- [15] Richhariya B, Tanveer M, Rashid AH. Diagnosis of Alzheimer's disease using universum support vector machine based recursive feature elimination (USVM-RFE). *Biomed Signal Process Contr*. 2020;59:101903. <https://doi.org/10.1016/j.bspc.2020.101903>
- [16] Wang S, Wang Y, Wang D, Yin Y, Wang Y, Jin Y. An improved random forest-based rule extraction method for breast cancer diagnosis. *Appl Soft Comput*. 2020;86:105941. <https://doi.org/10.1016/j.asoc.2019.105941>
- [17] Zhao H, Wang R, Lei Y, Liao WH, Cao H, Cao J. Severity level diagnosis of Parkinson's disease by ensemble K-nearest neighbor under imbalanced data. *Expert Syst Appl*. 2022;189:116113. <https://doi.org/10.1016/j.eswa.2021.116113>
- [18] Lesica NA, Mehta N, Manjaly JG, Deng L, Wilson BS, Zeng FG. Harnessing the power of artificial intelligence to transform hearing healthcare and research. *Nat Mach Intell*. 2021;3(10):840–9. <https://doi.org/10.1038/s42256-021-00394-z>

- [19] Murdoch WJ, Singh C, Kumbier K, Abbasi-Asl R, Yu B. Definitions, methods, and applications in interpretable machine learning. *Proc Natl Acad Sci USA*. 2019;116(44):22071–80. <https://doi.org/10.1073/pnas.1900654116>
- [20] Yuan H, Kang L, Li Y, Fan Z. Human-in-the-loop machine learning for healthcare: current progress and future opportunities in electronic health records. *Med Adv*. 2024;2(3):318–22. <https://doi.org/10.1002/med4.70>
- [21] Kulmanov M, Guzmán-Vega FJ, Duek Roggli P, Lane L, Arold ST, Hoehndorf R. Protein function prediction as approximate semantic entailment. *Nat Mach Intell*. 2024;6(2):220–8. <https://doi.org/10.1038/s42256-024-00795-w>
- [22] Oberste L, Heinzl A. User-centric explainability in healthcare: a knowledge-level perspective of informed machine learning. *IEEE Trans Artif Intell*. 2023;4(4):840–57. <https://doi.org/10.1109/TAI.2022.3227225>
- [23] Yuan H, Hong C, Jiang PT, Zhao G, Tran NTA, Xu X, et al. Clinical domain knowledge-derived template improves post hoc AI explanations in pneumothorax classification. *J Biomed Inform*. 2024;156:104673. <https://doi.org/10.1016/j.jbi.2024.104673>
- [24] Yang Z, Lin Y, Xu Y, Hu J, Dong S. Interpretable Disease Prediction via Path Reasoning over medical knowledge graphs and admission history. *Knowl Based Syst*. 2023;281:111082. <https://doi.org/10.1016/j.knosys.2023.111082>
- [25] Li R, Yin C, Yang S, Qian B, Zhang P. Marrying medical domain knowledge with deep learning on electronic health records: a deep visual analytics approach. *J Med Internet Res*. 2020;22(9):e20645. <https://doi.org/10.2196/20645>
- [26] Chen Y, Clayton EW, Novak LL, Anders S, Malin B. Human-centered design to address biases in artificial intelligence. *J Med Internet Res*. 2023;25:e43251. <https://doi.org/10.2196/43251>
- [27] Cuttillo CM, Sharma KR, Foschini L, Kundu S, MacKintosh M, Mandl KD, et al. Machine intelligence in healthcare—perspectives on trustworthiness, explainability, usability, and transparency. *NPJ Digit Med*. 2020;3(1):47. <https://doi.org/10.1038/s41746-020-0254-2>
- [28] Gennatas ED, Friedman JH, Ungar LH, Pirracchio R, Eaton E, Reichmann LG, et al. Expert-augmented machine learning. *Proc Natl Acad Sci USA*. 2020;117(9):4571–7. <https://doi.org/10.1073/pnas.1906831117>

How to cite this article: Yuan H. Clinical decision making: Evolving from the hypothetico-deductive model to knowledge-enhanced machine learning. *Med Adv*. 2024;2(4):375–9. <https://doi.org/10.1002/med4.83>